

7-Day Sleep Architecture Reset Protocol

Science-Based Circadian Recalibration for Deep, Restorative Sleep

For educational purposes only. Not medical advice. Consult a qualified sleep specialist or physician for your individual needs.

Sleep architecture describes the cyclical pattern of NREM (Stages N1, N2, N3) and REM sleep your brain cycles through roughly every 90 minutes. Disrupted architecture — caused by irregular schedules, light exposure, caffeine, alcohol, and stress — is the root cause of most non-apnoea sleep complaints. This 7-day protocol systematically recalibrates each architectural stage.

Rate each of the following 1 (never) to 5 (always) to identify your primary sleep disruption pattern:

- ☐ I take longer than 20 minutes to fall asleep most nights.
- ☐ I wake between 2–4 AM and struggle to return to sleep.
- ☐ I feel unrefreshed even after 7–8 hours in bed.
- ☐ I experience vivid, anxious dreams that disturb my sleep.
- ☐ I rely on an alarm clock to wake (without it, I would sleep significantly longer).
- ☐ I feel drowsy between 1–3 PM daily.

Total Score: __ / 30 ! Score 15+ = proceed with this full 7-day reset.

Ø=Ÿ, Science: Fixing a consistent wake time is the single most powerful intervention for circadian rhythm stabilisation. It regulates adenosine clearance and cortisol awakening response (CAR) timing.

- Choose a fixed wake time and commit to it for all 7 days — including weekends.
- Get 10+ minutes of natural outdoor light within 30 minutes of waking. This sets your circadian clock via retinal ipRGCs.
- Avoid all naps on Days 1 and 2 regardless of how tired you feel — this builds adenosine sleep pressure.
- Set your sleep window: count back 7.5 hours from your wake time (5 complete 90-min cycles).
- Place all screens outside the bedroom and charge them in another room.
- Set bedroom thermostat to 16–19 °C (60–66 °F) — the thermoregulatory drop triggers deep NREM N3 onset.
- Begin a 10-minute hot shower or bath 60–90 minutes before bed. The subsequent skin cooling accelerates sleep onset.
- Stop caffeine intake by 12:00 PM — caffeine's 5–6 hour half-life still occupies adenosine receptors at midnight.
- Take a 20-minute brisk walk before 5 PM. Moderate aerobic exercise increases N3 slow-wave sleep by up to 65%.
- Limit alcohol entirely — even 2 drinks suppress REM sleep for the entire first half of the night.

For educational purposes only. Not medical advice. Consult a qualified sleep specialist or physician for your individual needs.

- Begin a 10-minute "worry dump" journaling session 2 hours before bed — offload tomorrow's to-do list to paper.
- Practice box breathing for 5 minutes before sleep: inhale 4s, hold 4s, exhale 4s, hold 4s.
- Download f.lux or enable Night Shift/Night Mode on all screens from sunset onwards.
- Put on 60 BPM binaural beats or pink noise (not white noise — pink noise preferentially enhances N3 slow-wave).
- Read a physical book (not a screen) for 15 minutes in dim amber light as your final pre-sleep activity.
- Use a free sleep tracker (Apple Health, Google Fit, or a paper sleep diary) to log: time in bed, perceived sleep onset, wakes, and morning alertness 1–10.
- Reassess your original 6-item baseline — most people see 3–5 point improvements by Day 7.
- If you scored Stage N3 deficiency (unrefreshed sleep), add 400 mg magnesium glycinate 45 min before bed.
- If you scored REM disruption (vivid anxious dreams), add 300 mg ashwagandha with dinner.
- Commit to maintaining the fixed wake time permanently — this is the minimum effective dose for sustained sleep quality.

Time Before Bed	Action	Target Stage
3 hours	Final meal completed	Core body temp regulation
2 hours	No more alcohol or cannabis	REM protection
90 mins	Hot shower/bath	Thermoregulatory sleep onset
90 mins	Blue-light blocking glasses on	Melatonin onset
60 mins	Journaling & anxiety offload	Cortisol reduction
45 mins	Dimmed amber lighting only	Circadian cue
30 mins	Supplements if applicable	Biochemical preparation
15 mins	Physical book reading in bed	Behavioural conditioning
0 mins	Lights out, eyes closed	Sleep onset target

If you have not seen improvement after completing this protocol, consult a board-certified sleep medicine physician to rule out sleep apnoea (affecting 1 in 4 adults) or restless legs syndrome.